## **Author Index**

Abran, D., 389 Ananth, C. V., 260 Asensi, M., 406

Bard, H., 411 Barfield, W. D., 17 Bayet, T., 425 Blondel, B., 239 Buck, G. M., 31 Byrne, B. J., 461

Catalano, P. M., 181 Chemtob, S., 389 Claris, O., 439 Cnattingius, S., 25, 286 Cochrane, J., 296 Collins, J. H., 79 Conway, D. L., 225 Cotton, R. B., 387, 461 Craven, C., 36

Daval, J.-L., 416 De Halleux, V., 411 Della-Latta, P., 357 Demissie, K., 250 Devlieger, H., 425 Drysdale, S. M., 99 Dwyer, T., 296

Eugène, C., 425

Fifer, W. P., 89 Flood, P., 146 Flowers, L. M., 17 Foca, M. D., 332

Gagnon, C., 411 García-Sala, F., 406 Gelber, S. E., 346 Gibbs, R. S., 75 Goodman, S., 136 Grafe, M. R., 83 Graham, P. L., III, 322 Grojean, S., 416 Guignard, J.-P., 398 Gupta, A., 340 Haas, J. P., 367 Haataja, R., 450 Hallman, M., 450 Halpern, S. H., 122 Halsema, E. F. W., 447 Hankins, G., 1 Haxhiu, A., 432 Hou, X., 389 Hoyert, D. L., 3, 12

Iyasu, S., 17

Jaffe, R., 190 Johnson, C. D., 31 Joseph, K. S., 237, 250, 268

Kaminski, M., 239 Kinney, H. C., 83 Kirwan, J. P., 181 Kope, J. G., 447 Kramer, M. S., 237, 250 Kubin, C. J., 379

Lambe, M., 286 Landau, R., 109 Langer, O., 169, 196, 215 Leighton, B. L., 122 Lewis, D., 70 Lindstrom, D. P., 461 Lipillonne, A., 439 Liu, S., 268 Lloret, A., 406 Lombet, J., 425 Lubell, W. D., 389

McClintock, C., 51 Martin, J. A., 3, 12 Martin, R. J., 432 Martila, R., 450 Mazze, R. S., 171 Mellen, B. G., 461 Mhanna, M. J., 432 Miñana, J. B., 406 Muir, H., 99 Mulinare, J., 277 Myers, M. M., 89 Naudé, S., 425 Normand, S., 439

Olney, R. S., 277

Peri, K. G., 389 Picaud, J.-C., 439 Piper, J. M., 206 Platt, R. W., 306 Ponsonby, A.-L., 296

Quiniou, C., 389

Ratner, A. J., 346 Reygrobellet, B., 439 Rigo, J., 439 Rosenn, B. M., 210

Saade, G. R., 51 Saiman, L., 313, 315 Salle, B. L., 439 Samueloff, A., 232 Sastre, J., 406 Simpson, L. L., 42 Smiley, R., 9 Spiering, E. J. H., 447 Spong, C. Y., 1 Stephansson, O., 25

Tomashek, K. M., 17 Tressa, L. A., 367 Truttmann, A., 411

Van Sonderen, L., 447 Varma, D. R., 389 Vento, M., 406 Vert, P., 416 Viña, J., 406

Wapner, R. J., 70 Ward, K., 36 Wen, S. W., 260, 268 Willinger, M., 1 Wu, F., 357

Zakowski, M., 154

## Subject Index

A baumanii (Acinetobacter baumanii) infections, NICUacquired, 363, 383

AA (amino acid) content of PHPF, 442 Abdominal circumference (AC) measurement, determination of IUGR in GDM with, 193

Abortion, spontaneous, in GDM, prevention of, 199 Abscess, epidural, due to regional labor anesthesia, 154:155

AC (abdominal circumference) measurement, determination of IUGR in GDM with, 193 Accidents

placental, genetic and environmental links of tissue findings on peripartum fetal death due to, 38-39

see also Umbilical cord accidents, fetal death and Acetaminophen, 137, 150

Acetohexamide, 218

Acetylsalicylic acid, see Aspirin

Acinetobacter baumanii (A baumanii) infections, NICUacquired, 363, 383 Activated protein C resistance (APCR) in major inherited thrombophilias, stillbirths after 20 weeks and, 52, 54-55, 61, 62, 64

Adenovirus infections, NICU-acquired, 347, 352 Adrenocorticotropic hormone, 151 Adverse effect(s)

renal, of prostaglandin-synthesis inhibitors in premature infants, 398-405

see also Adverse effect(s) of CSE labor analgesia; Regional labor anesthesia, complication(s) of

Adverse effect(s) of CSE labor analgesia maternal hypotension, uterine hypertonus, and fetal bradycardia as, 116

PDPH as, see Postdural puncture headache pruritus as, 116-117

Adverse outcome

effects of prior pregnancy, on subsequent fetal death, population-based research on, 33-34 perinatal, smoking, overweight and, 291 see also specific conditions

AFLP (amplified fragment length polymorphism), molecular typing of microorganisms with, 362

Age, see Gestational age; Maternal age

Airway diseases

SP genes and, 456-458

see also Lung infections and diseases; Respiratory infections

Airway function, see Endogenous and exogenous nitric oxide, airway function and

Airway smooth muscle, NO and endogenous NO and contraction of, 433-434 see also Relaxation of airway smooth muscle, NO

Alleles, SP-A and SP-B, associated with risk for RDS, 453-454

Alpha (α)-hemolytic streptococcus infections, 318 Alphaprodine (á-prodine) in epidural labor analgesia, 131

Amantadine, 347, 351

Ambulation during labor, CSE labor analgesia and, 113-114

Ambulatory techniques of epidural labor analgesia, 106

Amide local anesthetics, see Bupivacaine; Levobupivacaine; Lidocaine; Ropivacaine

Amikacin, 381, 383, 385 Amino acid (AA) content of PHPF, 442

Aminoglycosides, see specific aminoglycosides Amoxicillin, 383, 384

Ampicillin, 328, 337, 342, 384

Amplification technology, nuclear, molecular typing of microorganisms with, 360-362

Amplified fragment length polymorphism (AFLP), molecular typing of microorganisms with, 362

Amplified polymorphic DNA (deoxyribonucleic acid), random, molecular typing of microorganisms with, 361

Análgesia, see Obstetric analgesia and anesthesia Anatomic considerations in CSE labor analgesia, 109-110 Anesthetic plan for nonobstetric surgery anesthesia in pregnancy, 143-144

Anesthetics

response to, in pregnancy, 138-139 see also specific anesthetic agents

Anomalies and malformations, see Congenital anomalies and malformations; Fetal growth abnormality(ies) in GDM

Antenatal conditions and interventions, see entries beginning with terms: Fetal

Antepartum fetal death

genetic and environmental links of tissue findings on, 39-40

see also Maternal medical condition(s), and risk of antepartum fetal death

Anterior spinal artery syndrome due to regional labor anesthesia, differential diagnosis between epidural abscess, epidural hemorrhage and, 155

Antibiotics, 77, 154

resistance to, see Microbial resistance to antibiotics as risk factors for NICU-acquired infections, 315, 317-319

see also specific antibiotics

Anticoagulation systems, natural, 51-52

Antimicrobial control programs in NICUs, 379-386 antimicrobial rotation in, 383

automated decision support programs for, 382 experience with, 383-385

guidelines and protocols for antimicrobial use in, 381-382

hospital formulary restrictions in, 380-381 optimal, 385 role of interdisciplinary antimicrobial

management teams in, 382

Antimicrobial management teams, role of interdisciplinary, 382

Antimicrobial rotation for NICU-acquired infections, 383

Antithrombin system, and inherited thrombophilias and antithrombin deficiency, 52, 53, 59, 60, 61, 64

limiting coagulation, 52

APCR (activated protein C resistance) in major inherited thrombophilias, stillbirths after 20 weeks and, 52, 54-55, 61, 62, 64

ASA, see Aspirin

Asphyxiated newborn infants resuscitated with pure oxygen, hypoxemia altering intra-erythrocytic redox status in, increasing GSSG and, 406-410

Aspiration test in CSE labor analgesia, 116 Aspirin (ASA; acetylsalicylic acid), 159

adverse renal effects of, 398-405 Association versus causation in epidural labor analgesia, 122-123

Astrocytes, hypertrophic, stillbirths after 20 weeks

and, 83, 86 Astrovirus infections, NICU-acquired, 347

Automated decision support programs for antimicrobial control, 382

Autonomic challenges, SIDS and, 90-91

Autonomic nervous system, development of, SIDS and, 92-93

Autosomal recessive disorders, stillbirths after 20 weeks and, 71

Aztreonam, 341

B cepacia (Burkholderia cepacia) infections, NICU-acquired, 383

Back pain, long-lasting, epidural labor analgesia and, 124, 132

Bacterial colonization

by CoNS, 322-323

hand, by P aeruginosa, 333-334

by S aureus, 324-325

Bacteriophage typing, 358

Bedding covering head, role of, in SIDS, 301-302

Benzodiazepines, teratogenicity of, 139-140

Beta-lactamases, see Extended spectrum β-lactamase Klebsiella pneumoniae infections, NICUacquired

Bilirubin, combined effects of hypoxia and, on forebrain neurons, 416-424

Biotyping of microorganisms, 358

Biparietal diameter, determination of IUGR in GDM with measurement of, 192

Birth(s), see Live births; Multiple births; Preterm births; Stillbirths

Birth weight

by gestational age in single pregnancies, trends in, 262-263, 266

as risk factor for NICU-acquired infections, 316; see also specific NICU-acquired infections see also Very low birth weight infants

Blood

epidural, 149, 161, 163

see also Inherited thrombophilias, stillbirths after 20 weeks and entries beginning with term: Blood and element: Hem-

Blood flow, umbilical

cessation of, in UCAs, stillbirths after 20 weeks and, 79

see also Disruption, umbilical blood flow

Blood gases in asphyxiated newborns resuscitated with pure oxygen or room air, 408

Blood loss, fetal, and disruption of umbilical blood flow, stillbirths after 20 weeks and, 79-80

Blood pressure, see Hypertensive disorders; Hypotension

Blood transfusions

effects of, on HbO<sub>2</sub> dissociation curve in very early preterm infants during first week of life, 411-415

NICU-acquired viral infections associated with, 353

Body-mass index (BMI), as risk factor for stillbirths after 20 weeks, 27, 28, 44, 47

Bone mineralization, nitrogen utilization and, VLBW infants on PHPF, 439-446

BPD, see Bronchopulmonary dysplasia

Bradycardia, fetal, due to CSE labor analgesia, 116 Brain damage, see Fetal brain damage Breastfeeding, epidural labor analgesia effects on, 124, 131-132

Bronchopulmonary dysplasia (BPD)

effects of exogenous NO on development of, 436-437

incidence of (1976-1997), 461-466

Bupivacaine, 112, 113, 116, 124-126, 129-131, 161-162

C albicans (Candida albicans) infections, NICU-acquired, 316-318, 363, 364

C diversus (Citrobacter diversus) infections, NICUacquired, 363

C freundii (Citrobacter freundii) infections, NICU-acquired, 341

C parapsilosis (Candida parapsilosis) infections, NICUacquired, 318, 363-364, 368, 374

C trachomatis (Chlamydia trachomatis) infections, 77 Caffeing, 147

Calicivirus infections, NICU-acquired, 347

Candida albicans (C albicans) infections, NICUacquired, 316-318, 363, 364

Candida infections, NICU-acquired, 317-319, 357, 379, 382

Candida parapsilosis (C parapsilosis) infections, NICUacquired, 318, 363-364, 368, 374

Candidate gene approach to RDS, 452-453

Capillary blood glucose, measurement of, in GDM, 172

Carbapenem, 337

Carbon dioxide (CO<sub>2</sub>) in nonobstetric surgery anesthesia, fetal effects of, 140

Cardiac arrest, regional labor anesthesia-related, 160 Cardiovascular system, physiological changes in, in pregnancy, 137

Case definitions in infection surveillance, defined, 368

Case finding in infection surveillance, defined, 368 Catheters

central venous, as risk factors for NICU-acquired infections, 315, 317, 324

testing of epidural, in CSE labor analgesia, 115-116

Causation, association versus, in epidural labor analgesia, 122-123

Cefazolin, 326, 381

Cefotan, 380

Cefotaxime, 341, 342, 381, 383, 384

Cefpodoxime, 341

Ceftazidime, 337, 340-343, 380, 384

Ceftriaxone, 341, 380

Cefuroxime, 342

Central venous catheters, as risk factors for NICUacquired infections, 315, 317, 324

Cephalosporins, see specific cephalosporins

Cerebral palsy, stillbirths after 20 weeks and, 83, 85 Cerebral white matter (perinatal telencephalic

leukoencephalopathy), stillbirths after 20 weeks and necrosis or gliosis of, 83-87

Cervical dilation with CSE labor analgesia, 114-115 Cesarean delivery epidural labor analgesia effects on rate of, 124, 126-128

see also Cesarean delivery of macrosomic fetus in GDM, vaginal versus

Cesarean delivery of macrosomic fetus in GDM, vaginal versus, 225-231

benefits of cesarean delivery in prevention of fetal injury, 226-227

fetal overgrowth and, see Fetal overgrowth in GDM

maternal morbidity associated with, 227-228

Cessation

of smoking by young women, trends in, 288 of umbilical cord blood flow in UCAs, stillbirths after 20 weeks and, 79

Chemical contamination, regional labor anesthesiarelated neurologic complications due to, 162

Child health, consequences of twin and/or triplet pregnancy on, 244-245

Chlamydia trachomatis (C trachomatis) infections, 77

Chloramphenicol, 335 Chlordiazepoxide, 139

2-Chloroprocaine, 162

Chlorpropamide, 218, 220

Cholestasis of pregnancy, and risk of antepartum fetal death, 47

Chorioamnionitis, stillbirths after 20 weeks and, 76,

Chromosomal analysis, molecular typing of microorganisms with, 360-361

Chronic renal disease, maternal, and risk of antepartum fetal death, 45-47

Cilastin, 380

Ciprofloxacin, 337, 343

Citrobacter diversus (C diversus) infections, NICUacquired, 363

Citrobacter freundii (C freundii) infections, NICU-acquired, 363

Citrobacter infections, NICU-acquired, 318, 334, 341, 363

Clavulanate, 337

Clavulanic acid, 343

Clindamycin, 381

Clinical decision-making on GDM, 172-179

Clinical presentation

of ESLB K pneumoniae infections, 342

of UCAs, as basis for evaluation of fetal risk for UCAs, 81

see also specific conditions

Clonidine, 102-103, 112, 130

CMV, see Cytomegalovirus infections

Coagulase-negative Staphylococcus aureus (S aureus) infections (CoNS), NICU-acquired, 317, 318, 322-323, 326, 328, 329

Codeine, 150

Cohorting during infection outbreaks in NICU, 370-973

Collectins, and innate immunity against lung infections, 456-458

Colonization, see Bacterial colonization

Collection, data, see Data collection on stillbirths after 20 weeks for national fetal death file Combined spinal-epidural (CSE) labor analgesia, 109-121

anatomic considerations in, 109-110

cervical dilation with, 114-115

epidural catheter testing in, 115-116

flexibility and rapid onset of, 113

maternal hypotension, uterine hypertonus, and fetal bradycardia with, 116

maternal satisfaction with, 113

motor block and ambulation with, 113-114 "needle through needle" technique of, 110-111

PDPH due to, see Postdural puncture headache pharmacology of, 111-113

pruritus due to, 116-117

safety of, 117

Comfort, maternal, epidural labor analgesia and, 123, 124

Compliance, respiratory system, with flow-pressure plot, 429-430

Complications, see Adverse effect(s); Complications, labor and delivery; Complications, maternal; Complications, pregnancy specific interventions and conditions

Complications, labor and delivery

perinatal death and (1995-1998), 22

smoking and overweight and, 290-291

see also Regional labor anesthesia, complication(s) of

Complications, maternal

at time of perinatal death (1995-1998), 22

see also Complications, labor and delivery; Complications, pregnancy; Neurologic complication(s), regional labor anesthesiarelated

Complications, pregnancy

of GDM, see Glucose thresholds, GDM complication(s) prevented with

severe, prevalence of inherited thrombophilias in women with, 64

smoking, overweight and, 289-290

Confined placental mosaicism (CPM), stillbirths after 20 weeks and, 71-72

Confirmation of infection outbreak in NICU, 369

Congenital anomalies and malformations

and fetal death at birth, genetic and environmental links in tissue findings on, 36-

see also Congenital anomalies and malformations, trends in fetal and neonatal mortality due to; Congenital anomalies and malformations in GDM; Neural tube defects, trends in prevalence of and specific conditions

Congenital anomalies and malformations, trends in fetal and neonatal mortality due to, 268-276

data sources on, 269-270

discussion on, 272-275 fetal mortality, 270-271

neonatal mortality, 271-272

Congenital anomalies and malformations in GDM glucose thresholds in prevention of, 199-201 oral hypoglycemic agents and, 218-219
CoNS (coagulase-negative Staphylococcus aureus (S

aureus) infections, NICU-acquired, 317, 318, 322-323, 326, 328, 329

Contamination, chemical, regional labor anesthesiarelated neurologic complications due to, 162

Context of vital statistics on stillbirths after 20 weeks, 13

Contractility, myometrial, THG113 effects on, 393-394

Contraction of airway smooth muscle, endogenous NO and 433-434

Cord accidents, see Umbilical cord accidents, fetal death and

Coronavirus infections, NICU-acquired, 352 Cortical vein thrombosis, as regional anesthesia complication, 159

Corticosteroids, 461, 464-465

Coumadin, 159

Coxsackie virus infections, NICU-acquired, 349

CPM (confined placental mosaicism), stillbirths after 20 weeks and, 71-72

Cranial nerve palsy in regional labor anesthesia, 163 CSE, see Combined spinal-epidural labor analgesia Cultural attitudes toward vital statistics on stillbirths after 20 weeks, 13

Cytogenetics at birth, stillbirths after 20 weeks and, 70-71

Cytomegalovirus (CMV) infections NICU-acquired, 346, 347, 353 stillbirths after 20 weeks and, 75, 77

Damage, see Fetal brain damage; Trauma and injury Data, national fetal death file

on causes of death, 9

collection of, see Data collection on stillbirths after 20 weeks for national fetal death file

filling gap on, see Population-based studies on fetal deaths

providers of, 14-15

quality of, 5-9

vital statistics as source of, see Vital statistics on stillbirths after 20 weeks

Data collection on stillbirths after 20 weeks for national fetal death file

instrument for, 13-14

recommendations on, 6

Data sources

on national fetal death file, see Vital statistics on stillbirths after 20 weeks

on trends in prevalence of folic acid fortification and multivitamin supplementation in NTDs, 277-269

Death, see Fetal mortality; Neonatal mortality Decision-making

on automated support programs for antimicrobial control in NICU, 382

clinical, on GDM, 172-179

Delayed preterm labor with THG113, 389-396

Delivery, see Cesarean delivery; Complications, labor and delivery; Delivery, effects of epidural labor analgesia on; Preterm delivery; Vaginal delivery

Delivery, effects of epidural labor analgesia on

on instrumental vaginal delivery, 124, 129 on rates of cesarean delivery, 124, 126-128

Demerol (meperidine), 129-131

Development, see Fetal overgrowth in GDM; Neurodevelopment and entries beginning with terms: Fetal growth

Dextran, 151, 159

Diabetes mellitus (DM), type 2, similarities between GDM and, oral hypoglycemic agents and, 215-216; see also Gestational diabetes mellitus

Diagnosis

differential, of epidural abscess, epidural hemorrhage, and anterior spinal artery syndrome due to regional labor anesthesia, 155

see also Imaging studies and specific conditions

Diazepam, 139

Differential diagnosis of epidural abscess, epidural hemorrhage, and anterior spinal artery syndrome due to regional labor anesthesia, 155

Dilation, cervical, with CSE labor analgesia, 114-115 Diphenhydramine, 116

Diseases associated with NICU-acquired infections with E faecalis infections, 327-328 with S aureus infections, 325-326 see also specific NICU-acquired infections

Disruption, umbilical blood flow fetal blood loss and, 79-80

intermittent, in UCAs, stillbirths after 20 weeks and, 79

Dissociation curve of HbO<sub>2</sub>, effects of blood transfusion on, in very early preterm infants during first week of life, 411-415

DM (diabetes mellitus), type 2, similarities between GDM and, oral hypoglycemic agents and, 215-216; see also Gestational diabetes mellitus

DNA (deoxyribonucleic acid), random amplified polymorphic, molecular typing of microorganisms with, 361

Dose

test, in epidural labor analgesia, controversies on, 104-105

see also specific pharmaceutical agents

Dystocia, fetal shoulder, fetal overgrowth in GDM and, 226-227

E aerogenes (Enterobacter aerogenes) infections, NICU-acquired, 363

E cloacae (Enterobacter cloacae) infections, NICUacquired, 318, 362, 363, 383, 384

E coli (Escherichia coli) infections, NICU-acquired, 318, 358, 362, 383, 384

E faecalis and E faecium infections, see: Enterococcus faecalis and Enterococcus faecium infections, NICU-acquired

Early neonatal mortality by state (1995-1998), 20 Echovirus infections, NICU-acquired, 349, 350 EFW (estimated fetal weight), determination of IUGR in GDM with, 193

Electrophoresis, pulsed-field gel, molecular typing of microorganisms with, 358-360

Endogenous and exogenous nitric oxide (NO), airway function and, 432-438

airway smooth muscle function, see Airway smooth muscle. NO and

effects of exogenous NO on neonatal lung injury and BPD development, 436-437

endogenous NO and airway smooth muscle contraction, 433-434

Endogenous flora, neonatal, acquisition of, 318, 319 Energy content of PHPF, 442, 444

Energy metabolism, neuronal, bilirubin effects on, 419

Enoxaparin, 159-160

Enteral nutrition, see Intravenous parenteral nutrition, NICU-acquired infections and

Enteric viral infections, NICU-acquired, 346-347
Enterobacter aerogenes (E aerogenes) infections, NICU-acquired, 363

Enterobacter cloacae (E cloacae) infections, NICU-acquired, 318, 341, 362, 363, 382-384

Enterobacter sazakii (E sazakii) infections, NICU-acquired, 374

Enterococcus faecalis (E faecalis) and Enterococcus faecium (E faecium) infections, NICU-acquired, 327-329

control of, 328-329

diseases associated with, 327-328

epidemiology of, 327

treatment of, 328

VRE, 322, 327, 328, 362, 379, 383, 384

Enteroviral infections, NICU-acquired, 349-350 Environment

cultural, of vital statistics collection on stillbirths after 20 weeks, 13

NICU, infection outbreaks and, 375

see also Tissue findings on stillbirths after 20 weeks, environmental and genetics links to EPB (epidural blood), 149, 151, 163

Epidemiology

of PDPH, 147-148

of RDS, 450-451

see also Epidemiology of NICU-acquired infections; Epidemiology of stillbirths after 20 weeks; Incidence; Perinatal epidemiology, trends in perinatal health and; Prevalence and specific conditions

Epidemiology of NICU-acquired infections enterococcal infections, 327 K pneumoniae infections, 340-341 P aeruginosa infections, 332-333 S aureus infections, 323-324

Epidemiology of stillbirths after 20 weeks, 25-30 of causes of stillbirths, 25 of risk factors for stillbirths, 26-29

role of gestational age in, 25-26

Epidural abscess, as regional anesthesia complication, 154-155

Epidural blood (EPB), 149, 151, 163 Epidural catheters, testing of, in CSE labor analgesia, 115-116

Epidural hematoma, as regional anesthesia complication, 159-160

Epidural hemorrhage in regional labor anesthesia, differential diagnosis of epidural abscess, anterior spinal artery syndrome and, 155

Epidural labor analgesia, 99-108 with ambulatory techniques, 106

with clonidine, 102-103

effects of, see Epidural labor analgesia, effect(s) of fluid preloading prior to, 105-106

with levobupivacaine, 99, 101-102

with neostigmine, 103

with PCEA, see Patient-controlled epidural analgesia

with ropivacaine, 99-101, 106, 113 test dose controversies in, 104-105

see also Combined spinal-epidural labor analgesia

Epidural labor analgesia, effect(s) of, 122-125 association versus causation and, 122-123 on breastfeeding, 124, 131-132

on delivery, see Delivery, effects of epidural labor analgesia on

on fetal and neonatal oxygenation, 123, 124 on length of labor, 123-126

long-lasting back pain as, 124, 132

on maternal comfort, 123, 124

on maternal temperature, 124, 129-131 urinary incontinence as, 124, 132-133

Epinephrine, 104-106, 114, 117, 130, 131, 159

Equipment, see Patient care equipment

Errors in gestational age, effects of, on interpretation of trends in fetal growth and gestational age-specific mortality, 306-311

ESBLs, see Extended spectrum β-lactamase infections, NICU-acquired

Escherichia coli (E coli), 318, 358, 362, 383, 384 Estimated fetal weight (EFW), determination of IUGR in GDM with, 193

Ethnic and socio-economic groups, SIDS rate reduction in different, 301

Ethnicity, see Race and ethnicity

Etiology

of PDPH, 146-147

see also specific conditions

Examination, neurologic, in regional labor anesthesia-related neurologic complications, 163-164

Exogenous nitric oxide, see Endogenous and exogenous nitric oxide, airway function and

Extended spectrum β-lactamase (ESBL) Klebsiella pneumoniae (K pneumoniae) infections, NICUacquired, 318, 319, 341-344

antimicrobial control programs for, 379

clinical presentation of, 342

epidemiology of, 341

molecular typing of ESBL K pneumoniae, 357, 363 prevention of, 343-344

treatment of, 342, 343

Extrinsic risk factors for NICU-acquired infections, 317-318

Factor V mutation in major inherited thrombophilias, stillbirths after 20 weeks and, 54-55, 60-64 Factor IX in major inherited thrombophilias, stillbirths after 20 weeks and, 57

Factor XI in major inherited thrombophilias, stillbirths after 20 weeks and, 57

Factor XIII polymorphism in major inherited thrombophilias, stillbirths after 20 weeks and,

Family visitation during NICU-acquired infection outbreaks, 374

Federal processing of vital statistics on stillbirths after 20 weeks, 15

Femur length (FL) measurement, determination of IUGR in GDM with, 193

Fentanyl, 97, 101, 102, 106, 112, 116, 125, 129, 130 Fetal anomalies and malformations, see Congenital anomalies and malformations and specific

Fetal blood loss, and disruption of umbilical blood flow, stillbirths after 20 weeks and, 79-80

Fetal bradycardia due to CSE labor analgesia, 116 Fetal brain damage

associated with stillbirths after 20 weeks, 83-88 damage to forebrain neuron due to combined effects of bilirubin and hypoxia, 416-424

Fetal characteristics

perinatal death and (1995-1998), 21 see also specific fetal characteristics

Fetal death, see Fetal mortality

Fetal effects

of nonobstetric surgery anesthesia, 139-142 see also entries beginning with terms: Maternal condition(s) and specific interventions

Fetal growth, trends in

effects of gestational age errors and their correction on interpretation of trends in gestational age-specific mortality and, 306-311 in single pregnancies, 260-267

Fetal growth abnormality(ies) in GDM, 190-195 IUGR as, 192-194

macrosomia as, 190-192; see also Fetal macrosomia in GDM

see also Fetal overgrowth in GDM

Fetal hemoglobin (HbF), effects of blood transfusion on, in very early preterm infants during first week of life, 411-415

Fetal injury due to fetal overgrowth, cesarean delivery preventing, 226-227

Fetal lung maturation in GDM, 206-209

Fetal macrosomia in GDM, 190-192

glucose thresholds and prevention of, 201-202 see also Fetal macrosomia in GDM,

pathophysiology of

Fetal macrosomia in GDM, pathophysiology of, 232-234

leptin in, 233-234

role of IGFs in, 233

role of insulin in, 232-233

Fetal monitoring in nonobstetric surgery anesthesia,

Fetal mortality

by state (1996-1998), 8

see also Abortion; Antepartum fetal death;

Congenital anomalies and malformations, trends in fetal and neonatal mortality due to: Data, national fetal death file; Late fetal deaths; Stillbirths and specific conditions

Fetal outcome

and anesthesia for nonobstetric surgery, 141-142 see also Fetal mortality specific interventions and

Fetal overgrowth in GDM, 225-227

detection of, with ultrasound, 228-230

pathology of, 225-226

shoulder dystocia and, 226-227

see also Cesarean delivery of macrosomic fetus in GDM, vaginal versus; Fetal macrosomia in

Fetal oxygenation, epidural labor analgesia effects

Fetal pathology, see Gestational diabetes mellitus. fetal pathophysiology in and specific conditions Fetal period of vulnerability to SIDS, 92

Fetal risk and risk factors

for fetal death, population-based research on, 33; see also Maternal medical condition(s), and risk of antepartum fetal death; Risk factor(s) for stillbirths after 20 weeks

for SIDS, 89-90

for UCAs, 81

Fetal shoulder dystocia, fetal overgrowth in GDM

Fibrinolysis system limiting coagulation, 52 First stage of labor, epidural analgesia effects on.

First week of life of very early premature infants, curve of HbO2 in, 411-415

FL (femur length) measurement, determination of IUGR in GDM with, 193

Flexibility of CSE labor analgesia, 113

acquisition of endogenous, 318, 319 patient-to-patient transmission of, 318-319

Flow-pressure plot, mechanical ventilation monitoring with, 425-431

discussion on, 430-431

in NICU, 428-429 output with, 426

physiological basis of, 427-428

system description, 426

validation of compliance and resistance

Fluid preloading, and epidural labor analgesia. 105-

Folate (folic acid) fortification for neural tube defects, trends in prevalence of, 243-244, 281

Formula, partially hydrolyzed preterm, nitrogen utilization and bone mineralization in VLBW infants fed, 439-446

Formulary restrictions, hospital, on antimicrobial use in NICU, 380-381

FP (prostaglandin<sub>2α</sub>) receptor antagonist, THG113 as, preterm labor delayed with, 389-396

Functional differentiation of lung, RDS and, 453

Gastrointestinal system, physiological changes in, in pregnancy, 137-138

GDM, see Gestational diabetes mellitus

Gel electrophoresis, pulse-field, molecular typing of microorganisms with, 358-360

Genetic and metabolic causes of stillbirths after 20 weeks, 70-74

cytogenetics at birth and, 70-71

Mendelian disorders, 71-73

see also Tissue findings on stillbirths after 20 weeks, environmental and genetics links to

Genetics, see Congenital anomalies and malformations; Genetic and metabolic causes of stillbirths after 20 weeks; Inherited thrombophilias, stillbirths after 20 weeks and; Molecular typing of microorganisms; Respiratory distress syndrome, genetics of; Twin(s)

Genotypes, association of SP-A and SP-B, with risk for RDS, 453-454

Genotypic molecular typing of microorganisms, 358 Gentamicin, 337, 341, 362, 381, 383-385

Germinal matrix hemorrhages (GMHs), stillbirths after 20 weeks and, 83-86

Gestational age

birth weight by, in single pregnancies, trends in, 262-263, 266

BPD and, 464

and effects of blood transfusion on HbO<sub>2</sub> dissociation, 413

effects of errors in, and their correction on interpretation of trends in gestational agespecific mortality, 306-311

of infants on PHPF, 443

see also Epidemiology of stillbirths after 20 weeks; Inherited thrombophilias, stillbirths after 20 weeks and; Tissue findings on stillbirths after 20 weeks, environmental and genetics links to and entries beginning with term: Preterm

Gestational age-specific mortality, effects of gestational age errors and their correction, on interpretation of trends in fetal growth and, 306-311

Gestational diabetes mellitus (GDM)

fetal pathophysiology in, see Gestational diabetes mellitus, fetal pathophysiology in

fetal testing in, 210-214

insulin sensitivity estimation and, 181-189

introduction to, 169-170 lung maturation in, 206-209

and risk of antepartum fetal death, 43-44, 47

see also Cesarean delivery of macrosomic fetus in GDM, vaginal versus; Fetal growth abnormality(ies) in GDM; Glucose thresholds, GDM complication(s) prevented with; Hyperglycemia in pregnancy, measurement of; Insulin sensitivity; Oral hypoglycemic agents in GDM

Gestational diabetes mellitus (GDM), fetal pathophysiology in, 232-236

in fetal growth, see Fetal growth abnormality(ies) in GDM

in oxygen delivery and iron, 234-235

GFR (glomerular filtration rate), effects of aspirin, indomethacin, and ibuprofen on, 401

Gliosis, cerebral white matter, stillbirths after 20 weeks and, 83-87

Glipizide, 220

Glomerular filtration rate (GFR), effects of aspirin, indomethacin, and ibuprofen on, 401

Glucose, see Gestational diabetes mellitus Glucose thresholds, GDM complication(s)

prevented with, 196-205 congenital malformations, 199-201 fetal macrosomia, 201-202 metabolic complications, 202

perinatal mortality, 197-199 respiratory complications, 202-203

spontaneous abortion, 199

Glutamate receptor antagonists, effects of, on bilirubin effects on forebrain neurons, 420

Glutathione, increasing oxidized, and hyperoxemia in asphyxiated infants resuscitated with pure oxygen altering intra-erythrocytic redox status, 406-410

Glyburide for GDM, 173, 176, 179, 217-220, 222 Glycosuria, measurement of, in GDM, 172

GMHs (germinal matrix hemorrhages), stillbirths after 20 weeks and, 83-86

Gram-negative bacteria molecular typing of, 362-363

see also specific bacteria

Gram-positive bacteria, see specific bacteria

Group B streptococcal infections, stillbirths after 20 weeks and, 77

Growth and development, see Fetal overgrowth in GDM; Neurodevelopment and entries beginning with terms: Fetal growth

GSSG (oxidized glutathione), hyperoxemia and increasing, altering intra-erythrocytic redox status in pure oxygen resuscitation of asphyxiated newborns, 406-410

Guidelines, see Recommendations and guidelines

H2 blockers, 318, 319

Hand colonization by P aeruginosa, 332-333

HAV (hepatitis A virus) infections, NICU-acquired, 348-349

Hb (hemoglobin), maternal concentrations of, stillbirths after 20 weeks and, 28-29

HbF (fetal hemoglobin), effects of blood transfusion on, in very early preterm infants during first week of life, 411-415

HbO<sub>2</sub> (hemoglobin oxygen), effects of blood transfusion on dissociation curve of, in very early preterm infants during first week of life, 411-415

HBV (hepatitis B virus) infections, transfusion-associated NICU-acquired, 353

HC (head circumference) measurement, determination of IUGR in GDM with, 192

HCV (hepatitis C virus) infections, transfusionassociated NICU-acquired, 353

Hcy (homocysteine), hyperhomocysteinemia and

475

5,10-MTHFR gene polymorphism in major inherited thrombophilias, 52, 55-57, 60-64

Head circumference (HC) measurement, determination of IUGR in GDM with, 192 Head covered with bedding, role of, in SIDS, 301-

302

Headache, see Postdural puncture headache Health, see Perinatal epidemiology, trends in perinatal health and

Healthy infants, acquisition of endogenous flora by, 318

Hearing loss due to regional labor anesthesia, 163 Hematologic system, physiological changes in, in pregnancy, 138

Hematoma due to regional labor anesthesia epidural, 159-160 subdural, 162

Hemoglobin (Hb), maternal concentrations of, stillbirths after 20 weeks and, 28-29

Hemoglobin oxygen (HbO<sub>2</sub>), effects of blood transfusion on dissociation curve of, in very early preterm infants during first week of life, 411-415

α (alpha)-Hemolytic streptococci, 318

Hemorrhage(s)

germinal matrix and intraventricular, stillbirths after 20 weeks and, 83-86

see also Hemorrhages due to regional labor anesthesia

Hemorrhages due to regional labor anesthesia epidural, differential diagnosis of epidural abscess, anterior spinal artery syndrome and, 155

subarachnoid, 159

Heparin, 159

Hepatitis A virus (HAV) infections, NICU-acquired, 348-349

Hepatitis B virus (HBV) infections, transfusionassociated NICU-acquired, 353

Hepatitis C virus (HCV) infections, transfusionassociated NICU-acquired, 353

Herpes simplex virus (HSV) infection NICU-acquired, 353

and regional anesthesia-related complications, 156 Heterogeneity of RDS and, 453

HIV (human immunodeficiency virus) infection transfusion-associated NICU-acquired, 353 see also HIV infections, maternal

HIV (human immunodeficiency virus) infections, maternal

and regional labor anesthesia-related neurologic complications, 163

stillbirths after 20 weeks and, 77

Homocysteine (Hcy), hyperhomocysteinemia and 5,1-MTHFR gene polymorphism in major inherited thrombophilias, 52, 55-57, 60-64

Hospital-acquired infections, see Neonatal intensive care unit-acquired infections

Hospital formulary restrictions on antimicrobial use, 380-381

HSV, see Herpes simplex virus infection Human immunodeficiency virus, see HIV infection Hyperbilirubinemia, forebrain neuron damage due to hypoxia and, 416-424

Hyperglycemia in pregnancy, measurement of, 171-180

clinical decision-making on, see Clinical decisionmaking on GDM

with continuous glucose monitoring, 176 glycosuria and capillary blood glucose

measurement, 172 with glycosylated hemoglobin and glycosylated protein measurement, 172-173

Hyperhomocysteinemia, 5,10-MTFHR polymorphism and, in major inherited thrombophilias, 52, 55-57, 60-64

Hyperoxemia in asphyxiated newborn infants resuscitated with pure oxygen, altering intraerythrocytic redox status, GSSG and, 406-410

Hyperoxia, effects of, on NO-mediated airway relaxant mechanisms, 434-436

Hypertensive disorders, maternal, and risk of antepartum fetal death, 42-43, 47

Hypertonus, uterine, due to CSE labor analgesia, 116

Hypertrophic astrocytes, stillbirths after 20 weeks and, 83, 86

Hypoglycemic agents, see Oral hypoglycemic agents in GDM

Hypotension, maternal, due to CSE labor analgesia, 116

Hypoxia, combined effects of bilirubin and, on forebrain neurons, 416-424

Ibuprofen, adverse renal effects of, 398-405 ICU (intensive care unit), see Neonatal intensive

Identification of infection outbreaks in NICU, 369 IGFs (insulin-like growth factors), role of, in fetal macrosomia in GDM, 233

Imaging studies

of fetal overgrowth with ultrasound, 228-230 of regional labor anesthesia related-neurologic complications in, 164

Imipenem, 343, 380

Immunity, innate, against lung infections, collectins and, 456-458

Immunologic deficiency, neonatal, as risk factor for NICU-acquired infections, 315-316

Improvement of vital statistics on stillbirths after 20 weeks, strategies for, 13

Incidence

of BPD (1976-1997), 461-466 UCA, 79, 81

of winter peaks of SIDS, attenuation of, 300 see also Epidemiology; Prevalence and specific

Incontinence, urinary, due to epidural labor analgesia, 124, 132-133

Indomethacin (INDO), adverse renal effects of, 398-405

Induction agents

teratogenicity of, 140 see also specific agents

Infants

health consequences of triplet pregnancies for, 239-240, 244-245

period of vulnerability of, to SIDS, 90

see also Preterm infants; Neonates; Sudden infant death syndrome, trends in; Sudden infant death syndrome, unexplained stillbirths and and entries beginning with terms. Neonatal, Perinatal

Infectious diseases

of lungs, see Lung infections and diseases stillbirths after 20 weeks due to, 75-78 see also Neonatal intensive care unit-acquired

infections

Infertility treatment, trends in multiple pregnancies and, 242-243, 245

Inflammatory lung diseases, SP genes and susceptibility to, 456

Influenza virus infections, NICU-acquired, 351-352 Information providers for vital statistics on stillbirths after 20 weeks, 14-15

Inhalation anesthetics

teratogenicity of, in nonobstetric surgery anesthesia, 139

see also specific inhalation anesthetics

Inherited thrombophilias, stillbirths after 20 weeks and, 51-69

major, see Major inherited thrombophilia(s), stillbirths after 20 weeks and

review of literature on, 57

systems limiting coagulation and, 51-52

Injury, see Trauma and injury

Innate immunity against lung infections, collectins and, 456-458

Inositol phosphate, effects of THG113 on  ${\rm PGF}_{2\alpha^+}$  induced production of, 392

Instrument for vital statistics collection on stillbirths after 20 weeks, 13-14

Instrumental vaginal delivery, epidural labor analgesia effects on rate of, 124, 129

Insulin, role of, in fetal macrosomia in GDM, 232-233

Insulin-like growth factors (IGFs), role of, in fetal macrosomia in GDM, 233

Insulin sensitivity (IS), estimation of, in pregnancy, 181-189

Insulin therapy of GDM, 215-218, 220

Intensive care unit, see Neonatal intensive care unit Interdisciplinary antimicrobial management teams, role of, 382

Intermittent disruption of blood flow in UCAs, stillbirths after 20 weeks and, 79

Intra-erythrocytic redox status, hypoxemia in pure oxygen resuscitation of asphyxiated newborns and alteration of, GSSG and, 406-410

Intralipids, intravenous, as risk factor for NICU-acquired infections, 317, 319

Intrathecal or intravascular injection, inadvertent, regional labor anesthesia-related neurologic complications due to, 162

Intrauterine growth retardation (IUGR) in GDM, 192-194

Intravascular devices, see Central venous catheters; Intravenous parenteral nutrition, NICUacquired infections and

Intravenous intralipids, as risk factor for NICUacquired infections, 317, 319

Intravenous parenteral nutrition, NICU-acquired infections and

infection outbreaks and handling of, 374 as risk factor for infections, 315, 317

Intraventricular hemorrhages (IVHs), stillbirths after 20 weeks and, 83-86

Intrinsic risk factors for NICU-acquired infections, 315-317

Invasive systems of continuous blood glucose monitoring in GDM, 176

Investigation of NICU-acquired infection outbreaks, see Outbreak of NICU-acquired infections Iron, fetal, in GDM, 234-235

IS (insulin sensitivity), estimation of, in pregnancy, 181-189

Ischemia, uterine, stillbirths after 20 weeks and, 80 Isolation of infected patients in NICU, 370-373

IUGR (intrauterine growth retardation) in GDM, 192-194

IVDs (intravascular devices), we Central venous catheters; Intravenous parenteral nutrition, NICU-acquired infections and

IVHs (intraventricular hemorrhages), stillbirths after 20 weeks and, 83-86

K aerogenes (Klebsiella aerogenes) infections, NICUacquired, 341

K oxytoca (Klebsiella oxytoca) infections, NICUacquired, 341

K planticola (Klebsiella planticola) infections, NICUacquired, 341

K pneumoniae, see. Klebsiella pneumoniae infections, NICU-acquired

K terrigena (Klebsiella terrigena) infections, NICUacquired, 341

Kanamycin, 333

Klebsiella aerogenes (K aerogenes) infections, NICU-acquired, 341

Klebsiella oxytoca (K oxytoca) infections, NICUacquired, 341

Klebsiella planticola (K planticola) infections, NICU-acquired, 341

Klebsiella pneumoniae (K pneumoniae) infections, NICU-acquired, 318, 334, 340-345

antimicrobial control programs and, 380, 381, 383, 384

characteristics of, 341-342

clinical presentation of, 342 epidemiology of, 340-341

ESBL, see Extended spectrum β-lactamase Klebsiella pneumoniae infections, NICUacquired

prevention of, 343-344

resistance of, to antibiotics, 319, 340-341, 343, 362, 379-381, 383-385

therapy of, 342-343

Klebsiella terrigena (K terrigena) infections, NICU-acquired, 341

L. monocytogenes (Listeria monocytogenes) infections, stillbirths after 20 weeks and, 77

Labor analgesia, see Obstetric analgesia and anesthesia

Labor and delivery, see Cesarean delivery; Complications, labor and delivery; Delivery, effects of epidural labor analgesia on; Preterm labor, prevention of; Vaginal delivery

Laboratory testing for neurologic complications of regional labor anesthesia, 164

β-Lactamases, see Extended spectrum β-lactamase Klebsiella pneumoniae infections, NICUacquired

Large for gestational age (LGA) births in single pregnancies, trends in, 266

Late fetal deaths

according to type of thrombophilia, 62 defined, 25

by state (1996-1998), 8

see also Perinatal mortality, late fetal death contribution to

Late neonatal mortality of less than 24 hours, by state (1996-1998), 8

Late-onset infections in NICU, 319

Length of labor, epidural labor analgesia effects on, 123-126

Leukoencephalopathy, perinatal telencephalic, stillbirths after 20 weeks and, 83-87

Levobupivacaine in obstetric analgesia and anesthesia, 99, 101-102

LGA (large for gestational age) births, in single pregnancies, trends in, 266

Lidocaine, 105, 140, 161-162

Linkage analysis of RDS, 452

Lipids, RDS and synthesis, transport, metabolism, and storage of, 452

Listeria monocytogenes (L. monocytogenes) infections, stillbirths after 20 weeks and, 77

Live births, number of perinatal death and, by year and race (1995-1998), 19

Local anesthetics

teratogenicity of, 139 see also specific anesthetics

Long-lasting back pain due to epidural labor analgesia, 124, 132

Lung(s)

RDS and functional differentiation of, 453 see also Bronchopulmonary dysplasia; Lung infections and diseases; Lung injury; Lung maturation

Lung infections and diseases collectins and innate immunity against, 456-458 SP genes and susceptibility to, 456

Lung injury, neonatal, effects of exogenous NO on, 436-437

Lung maturation in GDM, 206-209

Lupus erythematosus, maternal systemic, and risk of antepartum fetal death, 44-45, 47 M furfur (Malassezia furfur) infections, NICU-acquired, 317-319

M hominis (Mycoplasma hominis) infections, stillbirths after 20 weeks and, 77

Macronutrient content of PHPF, 442

Macrosomia, see Cesarean delivery of macrosomic fetus in GDM, vaginal versus; Fetal macrosomia in GDM

Magnesium sulfate in obstetric analgesia and anesthesia, 162

Magnitude of fetal deaths, population-based research on, 32

Major inherited thrombophilia(s), stillbirths after 20 weeks and, 52-57

activated protein C resistance in, 52, 54-55, 61, 62, 64

antithrombin deficiency in, 52, 53, 59, 60, 61, 64 factor V mutation in, 52, 54-55, 60-64

factor IX and XI in, 57

late fetal death according to type of, 62 PAI-1 gene polymorphism in, 57, 60

protein C deficiency as, 52-54, 60-62, 64 protein S deficiency as, 52, 54, 60-62, 64

protein 5 deficiency as, 52, 54, 60-62, 64 prothrombin G20210A mutation in, 52, 57, 60-62, 64

see also Major inherited thrombophilias, polymorphism in

Major inherited thrombophilias, polymorphism in 5,10-MTHFR gene polymorphism and hyperhomocysteinemia in, 52, 55-57, 60-64

of thrombomodulin gene C1418T, factor XIII gene, and PAI-1 gene, 57

see also Mutation in major inherited

Malassezia furfur (M furfur) infections, NICUacquired, 317-319

Maternal age mas determinant of multiple pregnancies, 241-242

as risk factor for stillbirths after 20 weeks, 26 Maternal body-mass index (BMI), as risk factor for stillbirths after 20 weeks, 27-28

Maternal characteristics perinatal death and (1995-1998), 21

see also specific maternal characteristics

Maternal comfort, epidural labor analgesia and,
123, 124

Maternal complications, see Complications, maternal Maternal health

consequences of twin and triplet pregnancies on, 239-240

see also Perinatal epidemiology, trends in perinatal health and entries beginning with terms: Maternal medical condition(s) and specific conditions

Maternal hemoglobin (Hb), stillbirths after 20 weeks and concentrations of, 28-29

Maternal human immunodeficiency virus infections, see HIV infections, maternal

Maternal hypotension due to CSE labor analgesia,

Maternal medical condition(s), and risk of antepartum fetal death, 42-50 cholestasis of pregnancy, 47 chronic renal disease, 45-47

GDM, 43-44, 47; see also Gestational diabetes mellitus

hypertensive disorders as, 42-43, 47

obesity as, 27, 28, 44, 47; see also Smoking and overweight during pregnancy, trends in SLE as, 44-45, 47

thyroid disorders as, 46-47

see also Maternal medical condition(s), stillbirths after 20 weeks and

Maternal medical condition(s), stillbirths after 20 weeks and

genetic and environmental links of tissue findings on stillbirths and, 40

at time of perinatal death (1995-1998), 22

see also Inherited thrombophilias, stillbirths after 20 weeks and; Maternal medical condition(s), and risk of antepartum fetal death

Maternal morbidity

associated with cesarean section delivery of macrosomic fetus, 227-22

see also entries beginning with terms: Maternal medical condition(s) and specific conditions

Maternal obesity, see Obesity, maternal

Maternal satisfaction with CSE labor analgesia, 113 Maternal temperature, epidural labor analgesia effects on, 124, 129-131

Maternal weight gain, see Weight gain, maternal Mechanical barrier to infections, 316

Mechanical ventilation, see Flow-pressure plot, mechanical ventilation monitoring with

Medical condition(s) severity of NICU patient, as risk for NICU-

acquired infections, 316-317 see also entries beginning with terms: Maternal medical condition(s) and specific conditions

Medication handling, and NICU-acquired infections, 317-318

Mendelian disorders, stillbirths after 20 weeks due to, 71-73

Meningitis, as regional anesthesia complication, 155-156

Meperidine (pethidine; Demerol), 129-131

Meprobamate, 139 Meropenem, 337, 343

Metabolic causes of stillbirths after 20 weeks, see Genetic and metabolic causes of stillbirths

after 20 weeks Metabolic stresses in fetus of GDM mother, 235 Metabolism, lipid, RDS and, 452

Methicillin-resistant Staphylococcus aureus (S aureus; MRSA) infections, NICU-acquired, 322-324, 326, 327, 362, 369, 379, 383, 384

Methohexital, 162

5,10-Methylenetetrahydrofolate reductase (5,10-MTHFR) gene, hyperhomocysteinemia and polymorphism of, in major inherited thrombophilias, 52, 55-57, 60-64

Metoclopramide, 129, 143 Metronidazole, 381

Microorganisms, see Molecular typing of

microorganisms specific microorganisms and infections

Microbial resistance to antibiotics

of CoNS infections, 322-323 of K pneumoniae infections, 319, 340-341, 343, 362, 379-381, 383-385

of P aeruginosa infections, 333, 335-337, 379-381, 383, 384

of S aureus infections, 322-324, 326, 327, 362, 369, 379, 383, 384

VRE infections, 322, 327, 328, 362, 379, 384

Mineralization, nitrogen utilization and bone, in VLBW infants on PHPF, 439-446

Model State Vital Statistics and Regulations (The Model Law), 3-5, 12

Molecular typing of microorganisms, 357-366 applications of, to specific organisms, 362-364 genotyping methods of, 358

with nonamplified technology, see Nonamplified molecular typing of microorganisms

with nucleic acid amplification technology, 360-

phenotypic methods of, 357-358

Monitoring

continuous glucose, in hyperglycemia, 176 fetal, in nonobstetric surgery anesthesia in pregnancy, 142

of health indicators concerning preterm births in twin pregnancies, 247

uterine, in anesthesia for nonobstetric surgery anesthesia in pregnancy, 142

see also Flow-pressure plot, mechanical ventilation monitoring with

Monoamniotic twins, stillborn, twin entanglement and, 80

Monochorionic twins, stillborn, brain damage and,

Morbidity, maternal, associated with cesarean versus vaginal delivery of macrosomic fetus in GDM, 227-228

Morphine, 113, 156

Morphology, neuronal nuclear, bilirubin effects on, 419, 421

Morphometric ratios, determination of IUGR in GDM with, 193

Mortality, see Fetal mortality; Neonatal mortality; Perinatal mortality and specific conditions

Mosaicism, confined placental, stillbirths after 20 weeks and, 71-72 Motor block with CSE labor analgesia, 113-114

MRSA (methicillin-resistant Staphylococcus aureus) infections, NICU-acquired, 322-324, 326, 327, 362, 369, 379, 383, 384

5,10-MTHFR (methylenetetrahydrofolate reductase) gene, polymorphism of, and hyperhomocysteinemia in major inherited thrombophilias, 55-57, 60-62, 64

Multidisciplinary team, investigation of NICUacquired infection outbreaks by, 369

Multifactorial disease, RDS as, 451-452

Multiple births

as risk factor for stillbirths after 20 weeks, 28-29

see also Multiple births, trends in

Multiple births, trends in, 239-249

in consequences on maternal and infant health, 239-240

in determinants of multiple pregnancies, 241-244 in triplet pregnancies, see Triplet pregnancies, trends in

in twin pregnancies, see Twin pregnancies, trends in

Multivitamin supplementation, trends in prevalence of, 243-244, 281-282

Muscle relaxants

teratogenicity of, 140
see also specific muscle relaxants

Mutation in major inherited thrombophilias APCR and, 52, 54-55, 61, 62, 64 factor V, 52, 54-55, 60-64 G20210A prothrombin, 52, 57, 60-62, 64 see also Polymorphism in major inherited

thrombophilias Mycoplasma hominis (M hominis) infections, stillbirths after 20 weeks and, 77

Myometrial contractility, effects of THG113 on, 393-394

Nalbuphine, 116 Naloxone, 116, 122

National Center for Health Statistics (NCHS), 4, 5, 9, 12, 18

National fetal death file, 3-11

data in, see Data, national fetal death file NCHS role and, 4, 5, 9 NVSS and, 45

National Institutes of Health Stillbirth Conference, see Vital statistics on stillbirths after 20 weeks

National Vital Statistics System (NVSS), 4-5, 12, 16 Natural anticoagulation systems, 51-52

NCHS (National Center for Health Statistics), 4, 5, 9, 12, 18

Necrosis, cerebral white matter, stillbirths after 20 weeks and, 83-87

"Needle through needle" technique of CSE labor analgesia, 110-111

Neonatal brain injury due to hyperbilirubinemia and hypoxia, 416-424

Neonatal characteristics, perinatal death and (1995-1998), 21

Neonatal health, consequences of triplet pregnancies on, 239-240; *see also* Perinatal epidemiology, trends in perinatal health and

Neonatal intensive care unit (NICU) flow-pressure system in, 428-429 see also Neonatal intensive care unit-acquired infections

Neonatal intensive care unit (NICU)-acquired infections

introduction to, 313-314

see also Antimicrobial control programs in NICUs; Enterococcus faecalis and Enterococcus faecium infections, NICU-acquired; Klebsiella pneumoniae infections, NICU-acquired; Molecular typing of microorganisms; Outbreak of NICU-acquired infections; Pseudomonas aeruginosa infections, NICU-acquired; Risk factor(s) for NICU-acquired infections; Staphylococcus aureus infections, NICU-acquired; Viral infections, NICU-acquired

Neonatal lung injury, effects of exogenous NO on, 436-437

Neonatal mortality

trends in gestational age-specific, effects of gestational age errors and their correction on interpretation of, 306-311

trends in preterm birth, obstetric intervention, stillbirths after 20 weeks and, 253-255

see also Congenital anomalies and malformations, trends in fetal and neonatal mortality due to; Neonatal mortality by states; Perinatal mortality; Stillbirths entries beginning with terms: Sudden infant death syndrome and specific conditions

Neonatal mortality by states early (1995-1998), 20 late (1996-1998), 8

Neonatal oxygenation, epidural labor analgesia effects on, 123, 124

Neonatal survival, and BPD incidence, 461-466 Neonates

hyperoxemia in pure oxygen resuscitation of asphyxiated, altering intra-erythrocytic redox status, GSSG and, 406-410

see also Infants and entries beginning with terms: Neonatal, Perinatal

Neostigmine, 103

Nerve trauma, regional labor anesthesia-related, 160-161

Nervous system, see Autonomic nervous system and entries beginning with element: Neur-

Neural tube defects (NTDs), trends in prevalence of, 277-285

data sources on, 277-279 discussion on, 282-283

of folic acid fortification, 243-244, 281

of multivitamin supplementation, 243-244, 281-282

and NTD rates, 279-281

Neurodevelopment, perinatal autonomic, SIDS and, 92-93

Neurologic complication(s), regional labor anesthesia-related, 156, 160-164 cardiac arrest and, 160

due to adverse effects of anesthetics, 161-162

due to chemical contamination, inadvertent intrathecal or intravascular injection, and subdural hematoma, 162

due to equipment problems, 161 hearing loss and cranial nerve palsy as, 163 in HIV-infected patients, 163 laboratory testing and imaging for, 164

nerve trauma as, 160-161

neurologic examination and testing for, 163-164 pneumocephalus as, 162-163

TRI and TNS as, 162

Neurologic examination and testing for neurologic complications of regional labor anesthesia,

Neurologic syndrome, transient, in regional labor anesthesia, 162

Neuron(s), forebrain, combined effects of bilirubin and hypoxia on, 416-424

Neuronal outcome, bilirubin and hypoxia effects on, 418

Neuronal viability and nuclear morphology, bilirubin effects on, 419, 421

Neuropathology associated with stillbirths after 20 weeks, 83-88

Newborns, see Neonates

NICU, see Neonatal intensive care unit

NIDDM (noninsulin-dependent diabetes mellitus; type 2 diabetes mellitus), similarities between GDM and, oral hypoglycemic agents and, 215-216; see also Gestational diabetes mellitus

NIH (National Institutes of Health) Stillbirth Conference, see Vital statistics on stillbirths after 20 weeks

Nitric oxide, see Endogenous and exogenous nitric oxide, airway function and

Nitrogen, bone mineralization and utilization of, in VLBW infants on PHPF, 439-446

Nitrous oxide (N<sub>2</sub>O), 139, 143

NO (nitric oxide), see Endogenous and exogenous nitric oxide, airway function and

Nonamplified molecular typing of microorganisms, 358-361

with chromosomal analysis, 360-361

with plasmid analysis, 361

with pulsed-field gel electrophoresis, 358-360 with restriction enzyme analysis, 359, 360

with ribotyping, 359-361

Noninsulin-dependent diabetes mellitus (NIDDM: type 2 diabetes mellitus), similarities between GDM and, oral hypoglycemic agents and, 215-216; see also Gestational diabetes mellitus

Noninvasive systems of continuous blood glucose monitoring in GDM, 176

Nonobstetric surgery in pregnancy, anesthesia for, 136-145

anesthetic plan for, 143-144 fetal considerations in, 139-142

physiological changes of pregnancy and, 136-139 recommendations and guidelines for, 142-144

Nonpatient-related factors in PDPH, 148-149 Nonsteroidal anti-inflammatory drugs (NSAIDs),

adverse renal effects of, 398-405

Nosocomial infections, see Neonatal intensive care unit-acquired infections

NSAIDs, see Nonsteroidal anti-inflammatory drugs NTDs, see Neural tube defects, trends in prevalence

Nuclear morphology, neuronal, bilirubin effects on, 419, 421

Nucleic acid amplification, molecular typing of microorganisms with, 360-362

Nutrition

of VLBW with PHPF, nitrogen utilization and bone mineralization in, 439-446 see also Intravenous parenteral nutrition, NICU-

acquired infections and; Supplementation NVSS (National Vital Statistics System), 4-5, 12, 16

Oo, see Oxygen

Obesity, maternal

and risk of antepartum fetal death, 27, 28, 44, 47 see also Smoking and overweight during pregnancy, trends in

Obstetric analgesia and anesthesia

introduction to, 97-98

see also Combined spinal-epidural labor analgesia; Epidural labor analgesia; Nonobstetric surgery in pregnancy, anesthesia for; Postdural puncture headache; Regional labor anesthesia, complication(s) of

Obstetric injuries, and regional anesthesia complications, 156-159

Obstetric intervention, see Preterm births, relationship between trends in obstetric intervention, stillbirths and

Onset of CSE labor analgesia, 113

Opioids

in obstetric analgesia, 97, 101, 109, 112, 115-117, 122, 123, 126, 128-132, 140, 143

see also specific opioids

Optimal antimicrobial control program, 385

Oral hypoglycemic agents in GDM, 215-224 clinical studies on, 220-221 fetal anomalies and, 218-219

placental transfer of, 219-220 rationale for, 216-218

similarities between type 2 diabetes mellitus and GDM and, 215-216

Oseltamivir, 347

Otitis media, SP genes and susceptibility to, 456 Outbreak of NICU-acquired infections, 367-378

case control studies of, 375 communicating findings on, 376

due to person-to-person transmission, 373-374

family visits during outbreak, 374 finding source of, 373

handling of medications and enteral feeds during, 374

identification of, 369

isolation and cohorting, 370-373

patient care equipment and, 374-375 state of NICU environment and, 375

surveillance of, 367-369

Outcome

neuronal, bilirubin and hypoxia effects on, 418 see also Fetal outcome; Neonatal mortality;

Neonatal survival; Neuronal outcome: Perinatal outcome; Pregnancy outcome and specific conditions

Output of flow-pressure system, 426

Overall risk factor for stillbirths after 20 weeks, plasma total homocysteine and (1967-1996), Overcrowding, understaffing and, as risk factors for NICU-acquired infections, 318

Overgrowth, fetal, see Fetal overgrowth in GDM Overweight, maternal, see Obesity, maternal Oxacillin, 384

Oxidative stress in pure oxygen resuscitation of asphyxiated newborns, hyperoxemia and, 406-410

Oxidized glutathione (GSSG), hyperoxemia and increasing, altering intra-erythrocytic redox status in pure oxygen resuscitation of asphyxiated newborns, 406-410

Oxygen (O<sub>2</sub>)

fetal delivery of, in GDM, 234-235

hemoglobin, effects of blood transfusion on dissociation curve of, in very early preterm infants during first week of life, 411-415

hyperoxemia in pure oxygen resuscitation of asphyxiated newborn infants altering intraerythrocytic redox status, GSSG and, 406-410 in nonobstetric surgery anesthesia, fetal effects of,

140

see also Oxygenation

Oxygenation

fetal and neonatal, epidural labor analgesia effects on, 123, 124

see also Flow-pressure plot, mechanical ventilation monitoring with

Oxytocin, 122, 125, 126

P<sub>50</sub>, effects of blood transfusion on, in very early premature infants, 411-415

PAI-1 (plasminogen activator inhibitor-1) gene polymorphism in major inherited thrombophilias and, 57

Palizumab, 347, 351

Palsy

cerebral, stillbirths after 20 weeks and, 83, 85 cranial nerve, in regional labor anesthesia, 163

Parainfluenza virus infections, NICU-acquired, 352 Parenteral nutrition, see Intravenous parenteral

nutrition, NICU-acquired infections and Parity, as risk factor for stillbirths after 20 weeks,

Partially hydrolyzed preterm formula (PHPF), nitrogen utilization and bone mineralization in VLBW infants on, 439-446

Parvovirus B19 infections, stillbirths after 20 weeks and, 75-77

Pathology and pathophysiology

of fetal overgrowth and, 225-226; see also Fetal macrosomia in GDM, pathophysiology of of PDPH, 146-147

see also Gestational diabetes mellitus, fetal pathophysiology in and specific conditions

Patient care equipment and infection outbreaks in NICU, 374-375 regional labor anesthesia-related neurologic

complications due to, 161 Patient-controlled epidural analgesia (PCEA), 103-104, 113, 118 Patient-to-patient transmission, see Person-to-person transmission in NICU

Patient-ventilator interaction, see Flow-pressure plot, mechanical ventilation monitoring with

PCEA (patient-controlled epidural analgesia), 103-104, 113, 118

PCR (polymerase chain reaction), see Repetitive polymerase chain reaction; Restrictive fragment length polymorphism-polymerase chain reaction

PDPH, see Postdural puncture headache Penicillin, 383-384

Perinatal autonomic nervous system development, SIDS and, 92-93

Perinatal epidemiology, trends in perinatal health and

introduction to, 237-238

see also Congenital anomalies and malformations, trends in fetal and neonatal mortality due to; Fetal growth, trends in; Multiple births, trends in; Neural tube defects, trends in prevalence of; Preterm births, relationship between trends in obstetric intervention, stillbirths and; Smoking and overweight during pregnancy, trends in; Sudden infant death syndrome, trends in

Perinatal mortality

in GDM, glucose thresholds and prevention of, 197-199

see also Perinatal mortality, late fetal death contribution to

Perinatal mortality, late fetal death contribution to (1995-1998), 17-24

and early neonatal mortality by state, 20 live births and, by year and race (1995-1998), 19 maternal, fetal, and neonatal characteristics and, 21

maternal medical conditions or complications at time of, 22

Perinatal outcome

trends in adverse, smoking, overweight and, 291 we also Perinatal mortality

Perinatal telencephalic leukoencephalopathy (PTL; cerebral white matter) necrosis or gliosis, stillbirths after 20 weeks and, 83-87

Period of vulnerability to SIDS fetal, 92

infant, 90

Peripartum fetal death due to placental and/or umbilical cord accidents, genetic and environmental links of tissue findings on, 38-

Peripheral nerve injuries in regional labor anesthesia, 157

Person-to-person transmission in NICU of flora, as risk factor for infections, 318-319 infection outbreaks and, 373-374 see also specific conditions

Pethidine (meperidine), 129-131

PFGE (pulsed-field gel electrophoresis), molecular typing of microorganisms with, 358-360 PG (prostaglandin)-synthesis inhibitors in premature infants, adverse renal effects of, 398-405

 $PGF_{2\alpha}$  (prostaglandin<sub>2 $\alpha$ </sub>; FP) receptor antagonist, THG113 as, preterm labor delayed with, 389-

Pharmacology of CSE labor analgesia, 111-113 Phenotypic methods of molecular typing of microorganisms, 357-358

PHPF (partially hydrolyzed preterm formula), nitrogen utilization and bone mineralization in VLBW infants on, 439-446

Physical properties of amide local anesthetics, 100 Physiological basis of flow-pressure plot, 427-428 Physiological changes in pregnancy, nonobstetric

surgery anesthesia and, 136-139

Piperacillin, 336, 337, 343, 381

Placental accidents, and peripartum fetal death, genetic and environmental links of tissue findings on, 38-39

Placental mosaicism, confined, stillbirths after 20 weeks and, 71-72

Placental transfer of oral hypoglycemic agents, 219-220

Plan, anesthetic, for nonobstetric anesthesia in pregnancy, 143-144

Plasma total homocysteine (Hcy), and overall risk factor for stillbirths after 20 weeks (1967-1996), 63

Plasmid analysis, molecular typing of microorganisms with, 361

Plasminogen activator inhibitor-1 (PAI-1) gene polymorphism in major inherited thrombophilias and, 57, 60

Pneumocephalus due to regional labor anesthesia, 162-163

Polymerase chain reaction, see Repetitive polymerase chain reaction; Restrictive fragment length polymorphism-polymerase chain reaction

Polymorphic DNA (deoxyribonucleic acid), random amplified, molecular typing of microorganisms with, 361

Polymorphism, see Polymorphism in major inherited thrombophilias; Surfactant protein gene polymorphism, predisposition to RDS and

Population-based studies

on RDS, 454

see also Population-based studies on fetal deaths Population-based studies on fetal deaths, 31-35

on magnitude and scope of fetal deaths, 32 of prior adverse pregnancy outcome effects on subsequent fetal death, 33-34

on probability of fetal deaths, 32-33

on risk factors for, 33; see also Maternal medical condition(s), and risk of antepartum fetal death; Risk factor(s) for stillbirths after 20 weeks

Postdural puncture headache (PDPH; spinal headache), 109, 117, 146-153 epidemiology of, 147-148 etiology and pathophysiology of, 146-147 nonpatient related factors in, 148-149 in pneumocephalus, 163 prevention of, 149-150 treatment of, 150-151

Predisposition to RDS, see Surfactant protein gene polymorphism, predisposition to RDS and

Pregnancy complications, see Complications, pregnancy

Pregnancy outcome

effects of prior adverse, on subsequent fetal death, 33-34

glucose profile and, 197; see also Glucose thresholds, GDM complication(s) prevented with

infectious diseases and, 76, 77 see also specific conditions

Prematurity, see Preterm births

Prenatal conditions and interventions, see entries beginning with terms: Fetal, Perinatal

Preterm births

after 20 weeks, genetic and environmental links of tissue findings on fetal death at birth due to, 36-38

see also Preterm births, relationship between trends in obstetric intervention, stillbirths and; Preterm births in twin pregnancies, trends in; Preterm delivery; Preterm infants

Preterm births, relationship between trends in obstetric intervention, stillbirths and, 250-259

trends in preterm birth, obstetric intervention, stillbirth, and infant mortality, 253-255

trends in preterm birth, stillbirth, and infant mortality, 252-253

trends in preterm birth and obstetric intervention, 252

Preterm births in twin pregnancies, trends in, 245-247

decreasing number of, 246-247 monitoring health indicators of, 247 prevention of, 245-246

Preterm delivery

induced by LPS, THG113 effects on, 394-395 of twin pregnancies, 244-245 see also Preterm births in twin pregnancies, trends in

Preterm formula, partially hydrolyzed, nitrogen utilization and bone mineralization in VLBW infants fed, 439-446

Preterm infants

adverse renal effects of prostaglandin-synthesis inhibitors in, 398-405

airway function in, see Endogenous and exogenous nitric oxide, airway function and BPD in, see Bronchopulmonary dysplasia

very early, during first week of life, effects of blood transfusion on dissociation curve of HbO<sub>2</sub> in, 411-415

VLBW infants on PHPF, nitrogen utilization and bone mineralization in, 439-446

see also Preterm births; Preterm delivery Preterm labor, prevention of in anesthesia for nonobstetric surgery, 140-141 with THG113, 389-396

Prevalence

of inherited thrombophilias in women with severe pregnancy complications, 64

see also Epidemiology; Incidence; Prevalence trends and specific conditions

Prevalence trends

in maternal overweight, 289-290

of risk factors for SIDS in different countries, 300 in smoking and smoking cessation by young

women, 287-288 see also Neural tube defects, trends in prevalence of

Prevention

of fetal injury, benefits of cesarean delivery in, 226-227

of GDM complications, see Glucose thresholds, GDM complication(s) prevented with

of NICU-acquired infections, see Prevention of NICU-acquired infections

of PDPH, 149-150

of preterm births in twin pregnancies, 245-246

of preterm labor, see Preterm labor

of regional anesthesia complications, 154 of SIDS, 297-300

of stillbirths after 20 weeks due to infectious diseases, 77

Prevention of NICU-acquired infections

of enterococcal infections, 328-329 of *K pneumoniae* infections, 343-344

of late-onset infections, 349-54

of S aureus infections, 324, 326-327

with vaccination, see Vaccination against NICUacquired infections

see also Antimicrobial control programs in NICUs Prior adverse pregnancy outcome, effects of, on

subsequent fetal death, 33-34

Probability of fetal deaths

population-based research on, 32-33 see also Risk factor(s) for stillbirths after 20 weeks

α-Prodine (alphaprodine) in epidural labor analgesia, 131

Prophylaxis, see Prevention

Prostaglandin (PG)-synthesis inhibitors in premature infants, adverse renal effects of, 398-405

Prostaglandin<sub>2 $\alpha$ </sub> (PGF<sub>2 $\alpha$ </sub>; FP) receptor antagonist, THG113 as, preterm labor delayed with, 389-396

Protein(s), neuronal, bilirubin effects on synthesis of, 420

Protein C deficiency in major inherited thrombophilias, stillbirths after 20 weeks and, 52-54, 62, 64

Protein C-protein S system limiting coagulation, 52 Protein C resistance, activated, factor V mutation and, in major inherited thrombophilias, stillbirths after 20 weeks and, 54-55, 60-62, 64

Protein S deficiency in major inherited thrombophilias, stillbirths after 20 weeks and, 52, 54, 62, 64

Prothrombin G20210A mutation in major inherited thrombophilias, 52, 57, 60-62, 64

Protocols, antimicrobial use, in NICU, 381-382

Providers of data for national fetal death file, 14-15 Pruritus due to CSE labor analgesia, 116-117

Pseudomonas aeruginosa (P aeruginosa) infections,

NICU-acquired, 332-339 antimicrobial control programs for, 380-384

categories of, 336-337

epidemiology of, 332-333

features of, 334-335

and hand colonization by P aeruginosa, 333-334

molecular typing in, 362

outbreaks of, 370, 374

person-to-person transmission of, 319

resistance of, to antibiotics, 333, 335-337, 379-381, 383, 384

therapy of, 337-338

PTL (perinatal telencephalic leukoencephalopathy), stillbirths after 20 weeks and, 83-87

Public health implications of SGA births, 266

Pulsed-field gel electrophoresis (PFGE), molecular typing of microorganisms with, 358-360

Puncture headache, see Postdural puncture headache

Pure oxygen (O<sub>2</sub>) resuscitation of asphyxiated newborns alteration of intra-erythrocytic redox status due to hyperoxemia in, increasing GSSG and, 406-410

Quality of national fetal death file data, 5-9 Quinolone, 337

Race and ethnicity

live births and perinatal deaths by year and (1995-1998), 19 and RDS, 454

Radicular irritation, transient, in regional labor anesthesia, 162

Random amplified polymorphic DNA (RAPD), molecular typing of microorganisms with, 361

RAR (room air resuscitation) of asphyxiated newborns, pure oxygen resuscitation compared with, 406-410

RDS, see Respiratory distress syndrome, genetics of REA (restriction enzyme analysis), molecular typing of microorganisms with, 359, 360

Recessive disorders, autosomal, stillbirths after 20 weeks and, 71

Recommendations and guidelines

AAP Task Force, on SIDS, 302-303

on antimicrobial use in NICU, 381-382

for nonobstetric surgery anesthesia in pregnancy, 142-144

on *S aureus* infection prevention in NICU, 324 Recurrence of post-20-week stillbirths, 28

Redox status, intra-erythrocytic, hypoxemia in pure oxygen resuscitation of asphyxiated newborns altering, increasing GSSG and, 406-410

Regional labor anesthesia, complication(s) of, 154-

cortical vein thrombosis as, 159 due to obstetric injuries, 156-159 epidural abscess as, 154-155 epidural hematoma as, 159-160 HSV infection as, 156 meningitis as, 155-156

neurologic, see Neurologic complication(s), regional labor anesthesia-related

prevention of, 154

subarachnoid hemorrhage as, 159 Relaxation of airway smooth muscle, NO and, 432-

and effects of hyperoxia or other pathologies on NO-mediated airway relaxant mechanisms, 434-436

exogenous NO and, 432-433

Renal disease, maternal chronic, and risk of antepartum fetal death, 45-47

Renal effects, adverse, of prostaglandin-synthesis inhibitors in premature infants, 398-405

Renal system, physiological changes in, in pregnancy, 138

Renovascular resistance (RVR), effects of aspirin, indomethacin, and ibuprofen on, 401

Repetitive polymerase chain reaction (PCR), molecular typing of microorganisms with, 361 Reporting requirements of Model State Vital

Statistics and Regulations, 3, 5 Resistance, measurement of respiratory system with

flow-pressure plot, 429-430 Respiratory complications in GDM, glucose thresholds and prevention of, 202-203

Respiratory distress syndrome (RDS), genetics of in twins, 447-449

see also Surfactant protein gene polymorphism, predisposition to RDS and

Respiratory infections NICU-acquired, 350-352

see also Lung infections and diseases

Respiratory syncytial virus (RSV) infections, NICUacquired, 350-351

Respiratory system

physiological changes in, in pregnancy, 137 see also Oxygenation entries beginning with terms: Airway, Respiratory

Restriction(s), hospital formulary, on antimicrobial use in NICU, 380-381

Restriction enzyme analysis (REA), molecular typing of microorganisms with, 359, 360

Restrictive fragment length polymorphismpolymerase chain reaction (RFLP-PCR), molecular typing of microorganisms with, 361-362

Resuscitation of asphyxiated newborn infants with pure oxygen, hypoxemia in altering intraerythrocytic redox status, increasing GSSG and, 406-410

Rett syndrome, stillbirths after 20 weeks and, 71 Revision of US Standard Report of Fetal death, 9-10 RFLP-PCR (restrictive fragment length

> polymorphism-polymerase chain reaction), molecular typing of microorganisms with, 361-362

Rhinovirus infections, NICU-acquired, 352 Ribotyping, molecular typing of microorganisms with, 359-361 Risk and risk factor(s) for BPD, 464

fetal, see Fetal risk and risk factors

prevalence of SIDS, in different countries, 300 see also Risk and risk factor(s) for NICU-acquired infections; Risk and risk factor(s) for RDS; Risk and risk factor(s) for stillbirths after 20 weeks

Risk and risk factor(s) for NICU-acquired infections, 315-321

acquisition of endogenous flora and, 318, 319 extrinsic, 317-318 intrinsic, 315-317

patient-to-patient transmission of flora and, 318-319

understaffing and overcrowding as, 318

Risk and risk factor(s) for RDS RDS-associated, 450-451

SP-A and SP-B alleles and genotypes associated with, 453-454

in twins, 448

Risk and risk factor(s) for stillbirths after 20 weeks epidemiology of, 26-29 multiple births as, 28-29

overall, plasma total homocysteine and, 63 population-based research on, 33

see also Maternal medical condition(s), and risk of antepartum fetal death; Maternal medical condition(s), stillbirths after 20 weeks and

Room air resuscitation (RAR) of asphyxiated newborns, pure oxygen resuscitation compared with, 406-410

Ropivacaine, 99-101, 106, 113

Rotation of antimicrobials in NICU, 383

Rotavirus (RV) infections, NICU-acquired, 347-348

RSV (respiratory syncytial virus) infections, NICUacquired, 347, 350-351

RV (rotavirus) infections, NICU-acquired, 347-348 RVR (renovascular resistance), effects of aspirin, indomethacin, and ibuprofen on, 401

S aureus infections, see. Staphylococcus aureus infections, NICU-acquired

S epidermidis (Staphylococcus epidermidis) infections, NICU-acquired, 318, 322

S maltophilia (Stenotrophomonas maltophilia) infections, NICU-acquired, 382, 383

S marcescens (Serratia marcescens) infections, NICUacquired, 341, 358, 362, 372

S pneumoniae (Streptococcus pneumoniae), NICU-acquired, 358

Safety of CSE labor analgesia, 117

Salmonella infection, NICU-acquired, 358

Satisfaction, maternal, with CSE labor analgesia, 113 Scope of fetal deaths, population-based research on, 32

SDM (Staged Diabetes Management), 176-179 Second stage of labor, epidural analgesia effects on,

Self-monitored blood glucose (SMBG), as basis for clinical decision-making on GDM, 172-179 Serotyping in NICU, 358 Serratia marcescens (S marcescens) infections, NICUacquired, 341, 358, 362, 372

Severe pregnancy complications, prevalence of inherited thrombophilias in women with, 64

Severity of NICU patient illness, as risk for NICUacquired infections, 316-317

SGA (small for gestational age) births in single pregnancies, trends in, 266

Shared features by SIDS and stillbirths, 93-94 Shigella infections, NICU-acquired, 358

Side sleeping, SIDS and, 301

SIDS, see Sudden infant death syndrome, trends in; Sudden infant death syndrome, unexplained stillbirths and

Single pregnancies, fetal growth in, trends in (1985-1998), 260-267

SLE (systemic lupus erythematosus), maternal, and risk of antepartum fetal death, 44-45, 47

Sleeping environments, unsafe, SIDS and, 303 Small for gestational age (SGA) births in single pregnancies, trends in, 266

SMBG (self-monitored blood glucose), as basis for clinical decision-making on GDM, 172-179

Smoking

as risk factor for stillbirths after 20 weeks, 27 see also Smoking and overweight during pregnancy, trends in

Smoking and overweight during pregnancy, trends in, 286-295

and adverse perinatal outcome, 291

overweight prevalence trends, 289-290 pregnancy and delivery complications and, 290-

smoking and smoking cessation prevalence in voung women, 287-288

Socio-economic factors

SIDS rate reduction and, 301 in stillbirths after 20 weeks, 28

Source of NICU-acquired infections, identifying, 373 SP, see Surfactant protein gene polymorphism,

predisposition to RDS and Spinal artery syndrome, anterior, differential diagnosis of epidural abscess, epidural hemorrhage and, in regional labor anesthesia. 155

Spinal-epidural labor analgesia, see Combined spinalepidural labor analgesia

Spinal headache, see Postdural puncture headache Spinal needle type, and PDPH incidence, 148-149 Spontaneous abortion in GDM, glucose thresholds

and prevention of, 199

Staged Diabetes Management (SDM), 176-179
Staphylococcus aureus (S aureus) infections, NICU-acquired, 322-327

coagulase-negative, 317, 318, 322-323, 326, 328, 329

colonization by S aureus, 324-325

diseases due to, 325-326

epidemiology of, 323-324

molecular typing of, 357, 358, 362

MRSA infections, 322-324, 326, 327, 362, 369, 379, 383, 384

prevention of, 324, 326-327 treatment of, 326

Staphylococcus epidermidis (S epidermidis) infections, NICU-acquired, 318, 322

States

late fetal mortality by (1996-1998), 8

processing of vital statistics on still births after 20 weeks by, 15

see also Neonatal mortality by states

Statistics, see Vital statistics on stillbirths after 20 weeks

Stenotrophomonas maltophilia (S maltophilia) infections, NICU-acquired, 382, 383

Steroids, 159

Stillbirths

defined, 1, 3, 5

due to infectious diseases, 75-78

introduction to, 1-2

neuropathology associated with, 83-88

see also Epidemiology of stillbirths after 20 weeks; Genetic and metabolic causes of stillbirths after 20 weeks; Inherited thrombophilias, stillbirths after 20 weeks and; Late fetal deaths; Maternal medical condition(s), and risk of antepartum fetal death; National fetal death file; Perinatal epidemiology, trends in perinatal health and; Perinatal mortality, late fetal death contribution to; Population-based studies on fetal deaths; Preterm births, relationship between trends in obstetric intervention, stillbirths and; Sudden infant death syndrome, unexplained stillbirths and; Tissue findings on stillbirths after 20 weeks, environmental and genetics links to; Twin(s), stillborn, after 20 weeks; Vital statistics on stillbirths after 20 weeks

Storage, lipid, RDS and, 452

Streptococcal infections

group B, stillbirths after 20 weeks and, 77 NICU-acquired alpha-hemolytic, 318

Streptococcus pneumoniae (S pneumoniae) infections, NICU-acquired, 358

Stress, oxidative, in pure oxygen resuscitation of asphyxiated newborns, hyperoxemia and, 406-410

Subarachnoid hemorrhage, as regional anesthesia complication, 159

Subdural hematoma due to regional labor anesthesia, 162

Subsequent fetal death, effects of prior adverse pregnancy outcome on, 33-34

Succinylcholine, 139

Sudden infant death syndrome (SIDS), trends in, 296-305

AAP Task Force recommendations on (1998-1999), 302-303

data sources on, 296-297

features of, in postintervention era, 300-302 prevention of, 297-300

trends since 1992, 300

Sudden infant death syndrome (SIDS), unexplained stillbirths and, 89-96

autonomic challenges in, 90-91
fetal risk factors for, 89-90
perinatal autonomic development and, 92-93
period of vulnerability to, see Period of
vulnerability to SIDS
shared features between, 93-94
and SIDS relationship to stillbirths, 91-92
Sufentanil, 97, 102, 106, 112, 114, 116, 117, 130
Sulbactam, 341, 343, 384
Sulfamethoxazole, 341
Sulfonylureas in GDM, 217-221
Sumatriptan, 147, 150
Supplementation

multivitamin, prevalence trends in, 243-244, 281-282 vitamin A, in BPD, 461, 464

Surfactant protein (SP) gene polymorphism, predisposition to RDS and, 450-460 association of SP-A and SP-B alleles and genotypes

with risk for RDS, 453-454 candidate gene approach to, 452-453

epidemiology of RDS and risk factors associated with, 450-451

heterogeneity of RDS and, 453 linkage analysis of, 452

and other lung and airway diseases, 456-458 and RDS as multifactorial disease, 451-452

Surfactant therapy, 461, 464 Survival, neonatal, BPD incidence and, 461-466 Syncytial virus, NICU-acquired respiratory, 347, 350-

351 Syphilis, untreated, pregnancy outcome with, 76 System(s) limiting coagulation, 51-52

Systemic lupus erythematosus (SLE), maternal, and risk of antepartum fetal death, 44-45, 47

Tazobactam, 337, 341, 343

Telencephalic leukoencephalopathy (cerebral white matter) necrosis or gliosis, perinatal, stillbirths after 20 weeks and, 83-87

Temperature, maternal, epidural labor analgesia effects on, 124, 129-131

Teratogenicity of anesthetics, 139-140 of benzodiazepines, 139-140 defined, 139

of inhalation anesthetics, 139 of opioids, muscle relaxants, induction agents, and local anesthetics, 140

Test dose in epidural labor analgesia, 104-105 TF (tissue factor) pathway inhibitor system limiting coagulation, 51-52

Theophylline, 147 Therapy

of BPD, 461, 464-465

of GDM, see Clinical decision-making on GDM; Oral hypoglycemic agents in GDM

infertility, trends in multiple pregnancies and, 242-243, 245

of NICU-acquired infections, see Therapy of NICU-acquired infections of PDPH, 150-151

preventive, see Prevention

see also specific drugs, therapeutic modalities, and conditions

Therapy of NICU-acquired infections of enterococcal infections, 328

of *K pneumoniae* infections, 342-343 of *P aeruginosa* infections, 337-338 of *S aureus* infections, 326

THG113, as  $PGF_{2\alpha}$  receptor antagonist, preterm labor delayed with, 389-396

Thiopental, 138, 162

Thromboembolism, venous, see Inherited thrombophilias, stillbirths after 20 weeks and Thrombomodulin gene C1418T, polymorphism of,

in major inherited thrombophilias, 57 Thrombophilias, see Inherited thrombophilias, stillbirths after 20 weeks and

Thrombosis, cortical vein, as regional anesthesia complication, 159

Thyroid disorders, maternal, and risk of antepartum fetal death, 46-47

Ticarcillin, 336, 337

Timing of anesthesia for nonobstetric surgery anesthesia in pregnancy, 142

Tissue factor (TF) pathway inhibitor system limiting coagulation, 51-52

Tissue findings on stillbirths after 20 weeks, environmental and genetics links to, 36-41 accidents of placenta and umbilical cord and fetal

death in peripartum period, 38-39 in antepartum fetal death, 39-40

role of prematurity and congenital malformations in fetal death at birth, 36-38

on unidentified stillbirths, 40-41

see also Maternal medical condition(s), stillbirths after 20 weeks and

TNS (transient neurologic syndrome) in regional labor anesthesia, 162

Tobramycin, 337, 380, 383-384 Tolbutamide, 220

Total homocysteine (Hcy), and overall risk factor for stillbirths after 20 weeks (1967-1996), 63

Toxoplasmosis, stillbirths after 20 weeks and, 75-77 Transfusion, see Blood transfusions

Transient neurologic syndrome (TNS) in regional

labor anesthesia, 162 Transient radicular irritation (TRI) in regional

labor anesthesia, 162 Transmission of infections, see Person-to-person transmission in NICU

Transport, lipid, RDS and, 452

Trauma and injury

brain, see Fetal brain damage

neonatal lung, effects of exogenous NO on, 436-437

nerve, regional labor anesthesia-related, 160-161 obstetric, and regional anesthesia complications, 156-159

Treatment, see Therapy

Trends in perinatal health, see Perinatal epidemiology, trends in perinatal health and

TRI (transient radicular irritation) in regional labor anesthesia, 162

Trimethoprim, 335, 341

Triplet pregnancies, trends in, 239-241 in consequences on maternal health, infant

health, and child health, 239-240, 244-245 in number, 240, 241

Twin(s)

genetics of respiratory distress syndrome in, 447-449, 454-455

see also Twin(s), stillborn, after 20 weeks; Twin pregnancies

Twin(s), stillborn, after 20 weeks monoamniotic, due to entanglement, 80 monochorionic, brain damage and, 86

Twin pregnancies RDS in, 454-455

see also Twin pregnancies, trends in

Twin pregnancies, trends in, 239-241 in consequences for child health, 244-245 in consequences for maternal and infant health,

in number of twin pregnancies, 239, 241, 244-245 see also Preterm births in twin pregnancies, trends

Type 2 diabetes mellitus (noninsulin-dependent diabetes mellitus), similarities between GDM and, oral hypoglycemic agents and, 215-216; see also Gestational diabetes mellitus

Typing, molecular, see Molecular typing of microorganisms

U urealyticum (Ureaplasma urealyticum) infections, stillbirths after 20 weeks and, 77

UCAs, see Umbilical cord accidents, fetal death and Ultrasound, fetal overgrowth detection with, 228-230 Umbilical blood flow, see Blood flow, umbilical Umbilical cord abnormalities, stillbirths after 20

weeks and, 80 Umbilical cord accidents (UCAs), fetal death and, 79-82

clinical signs of, in evaluation of at-risk fetus, 81 genetic and environmental links of tissue findings on peripartum fetal death due to, 38-39 incidence of, 79

possible mechanisms of, 79-80

Understaffing, overcrowding and, as risk factors for NICU-acquired infections, 318

Unidentified fetal death, genetic and environmental links of tissue findings on, 40-41

Unsafe sleeping environments, SIDS and, 303 Untreated syphilis, pregnancy outcome with, 76 Ureaplasma urealyticum (U urealyticum) infections, stillbirths after 20 weeks and, 77

Urinary incontinence, epidural labor analgesia and, 124, 132-133

Urine flow rate (UV), effects of aspirin, indomethacin, and ibuprofen on, 401

US Standard Report of Fetal Death, 3, 12 revision of, 9-10

Uterine hypertonus due to CSE labor analgesia, 116 Uterine ischemia, stillbirths after 20 weeks and, 80 Uterine monitoring in anesthesia for nonobstetric surgery anesthesia in pregnancy, 142

UV (urine flow rate), effects of aspirin, indomethacin, and ibuprofen on, 401

Vaccination against NICU-acquired infections influenza, 351-352 VZV, 353

Vaginal delivery

instrumental, epidural labor analgesia effects on rate of, 124, 129

see also Cesarean delivery of macrosomic fetus in GDM, vaginal versus

Validation of respiratory system compliance with flow-pressure plot, 429-430

Vancomycin-resistant microorganisms, 319, 326, 337 VRE, 322, 327, 328, 379, 383, 384

Varicella zoster virus (VZV) infections, NICUacquired, 352-353

Vasculitis, stillbirths after 20 weeks and, 76 Vasomotor response to PGF<sub>2α</sub>, effects of THG113

on, 392-393 Velocity of fetal growth, measurement of, determination of IUGR in GDM with, 193-

Venous thromboembolism, see Inherited thrombophilias, stillbirths after 20 weeks and

Ventilation, see Flow-pressure plot, mechanical ventilation monitoring with; Oxygenation

Very early preterm infants, effects of blood transfusion on HbO, dissociation curve in, during first week of life, 411-415

Very low birth weight (VLBW) infants on PHPF, nitrogen utilization and bone mineralization in, 439-446 see also Preterm infants

Viability, neuronal, bilirubin effects on, 419, 421

Viral infections, NICU-acquired, 346-356 due to CMV, 346, 347, 353 due to enteroviruses, 347, 349-350

due to rotavirus, 347-348 enteric, 346-347 hepatitis A, 348-349

herpes simplex, 347, 353 respiratory, 350-352

transfusion-associated, 353 varicella zoster, 347, 352-353

Virulence of CoNS, 322

of K pneumoniae, 342

Vital statistics on stillbirths after 20 weeks, 12-16 background to, 12

culture and context of, 13

data collection for, see Data collection on stillbirths after 20 weeks for national fetal death file

information providers on, 14-15 potential uses of, 15-16

state and federal processing of, 15 strategies for improvement of, 13

Vitamin A supplementation in BPD, 461, 464 VRE (vancomycin-resistant enterococcal) infections, NICU-acquired, 322, 327, 328, 379, 383, 384

- VTE (venous thromboembolism), see Inherited thrombophilias, stillbirths after 20 weeks and Vulnerability to SIDS, see Period of vulnerability to
- SIDS
- VZV (varicella zoster virus) infections, NICU-acquired, 352-353
- Weight
  - bone mineralization and nitrogen utilization in VLBW infants on PHPF, 439-446; see also Preterm infants
  - see also Birth weight; Obesity, maternal; Weight gain, maternal

- Weight gain, maternal
  - as risk factor for stillbirths after 20 weeks, 28 see also Smoking and overweight during
    - pregnancy, trends in
- Winter peaks of SIDS, trends in incidence of, 300
- X-linked disorders, stillbirths after 20 weeks and, 71
- Yeast, molecular typing of, 363-363
- Young women, trends in smoking and smoking cessation by, 287-288

